

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 9/96, A61K 47/26	A1	(11) International Publication Number: WO 91/18091 (43) International Publication Date: 28 November 1991 (28.11.91)
(21) International Application Number: PCT/GB91/00759 (22) International Filing Date: 14 May 1991 (14.05.91) (30) Priority data: 9010742.6 14 May 1990 (14.05.90) GB (71) Applicant (for all designated States except US): QUADRANT HOLDINGS CAMBRIDGE LIMITED [GB/GB]; Cambridge Research Laboratories, 181A Huntingdon Road, Cambridge CB3 0DJ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : ROSER, Bruce, Joseph [AU/GB]; The Old Vicarage, Church Lane, Balsham, Cambridgeshire CB1 6OS (GB). COLACO, Camilo [GB/GB]; 107 Foster Road, Trumpington, Cambridgeshire CB2 2JN (GB).		(74) Agent: ABLEWHITE, Alan, James; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US. Published <i>With international search report.</i>
(54) Title: STABILIZATION OF BIOLOGICAL MACROMOLECULAR SUBSTANCES AND OTHER ORGANIC COMPOUNDS (57) Abstract A method of preserving delicate biological substances or organic compounds (a) in a dry state and/or (b) at elevated temperatures and/or (c) under irradiation comprises incorporating in a system containing the said substances or compounds a sugar or a sugar derivative selected from (i) a non-reducing glycoside of a polyhydroxy compound selected from sugar alcohols and other straight chain polyalcohols, or (ii) a non-reducing oligosaccharide selected from raffinose, stachyose and melezitose.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

Stabilization of Biological Macromolecular
Substances and other organic compounds

This invention relates to the stabilization of biological macromolecular substances, in particular proteins and the like, such as enzymes, antibodies, fluorescent agents, vaccines etc., under dry and/or hot and/or irradiated conditions.

Our earlier patent (GB 2,187,171, U.S. 4,891,319) and applications (PCT/GB89/00047, PCT/GB89/00093) disclosed and claimed the use of the non-reducing sugar trehalose to preserve biological compounds while being dried at ambient temperature. An additional advantage of that process is that the compounds are not only stabilized during the drying procedure, but are also stabilized against long-term storage at relatively high ambient temperatures. Our earlier application (EP-A-0415567) disclosed the stabilisation also of low-molecular weight compounds such as antibiotics, chelating agents and nucleosides. Other compounds needing protection include vitamins.

Trehalose is a highly effective agent for this purpose and, in our subsequent research, has effectively provided a standard against which any other stabilizing agent must be compared. However, suitably pure

trehalose is not readily available in large quantities. There is therefore a need for alternative stabilizing agents. Sugars in general are of limited use. For most purposes reducing sugars do not provide the degree of stabilization required, especially during the drying procedure at a relatively high ambient temperature. Furthermore, reducing sugars are found to be of very little help in long-term stabilization and, in some cases can be highly deleterious. Other apparently non-reducing sugars, such as sucrose, again have limited utility and, even if they provide good stabilization during the drying step, they again provide very inferior long-term stabilization.

A chance observation that, under unusual circumstances, trehalose will crystallize on drying and that biological materials associated with the crystalline trehalose lacked protection, led to a search for compounds which could be dried without crystallizing, which were non-reducing but which were polyhydroxy compounds capable of replicating the effect of trehalose. We have now found that one particular type of carbohydrate material fulfills these requirements and is extremely effective in stabilizing biological materials both during the drying step and for long term storage.

The compounds in question are non-reducing glycosides of polyhydroxy compounds selected from sugar

SUBSTITUTE SHEET

alcohols and other straight chain polyalcohols. Other useful compounds include raffinose, stachyose and melezitose.

The sugar alcohol glycosides are preferably monoglycosides, in particular the compounds obtained by reduction of disaccharides such as lactose and maltose. The glycosidic group is preferably a glucoside or a galactoside and the sugar alcohol is preferably sorbitol (glucitol).

Particularly preferred compounds are maltitol (4-O- β -D-glucopyranosyl-D-glucitol), lactitol (4-O- β -D-galactopyranosyl-D-glucitol) and the two isomers of palatinit, the sugar alcohol derived from isomaltulose (palatinose) (6- α -D-glucopyranosyl-mannitol and 6- α -D-glucopyranosyl-sorbitol).

In comparative tests we have shown that these compounds are surprisingly very effective in contrast to other sugar alcohols and other sugars, whether reducing or not.

An example of the beneficial effect of the compounds according to this invention is as follows:-

Molecular biology techniques which form the basis for the technology of genetic engineering depend on the use of highly specific reagents. An important class of such reagents are the so called restriction enzymes. These proteins, largely derived from bacterial sources, have the property of recognising particular sequences of nucleotides in the DNA chain where they cause scission of the inter nucleotide bonds and cut the DNA into precise and reproducible fragments. These enzymes vary considerably in stability. Some are stable even at elevated temperature while others are among the most fragile biological reagents known. As a consequence, these enzymes are difficult and expensive to store, ship and handle. Routinely, these reagents are shipped in small aliquots of a few micrograms packed in solid CO₂ in large foam-polystyrene containers and transported by express services usually by air. This adds considerably to their cost. On receipt they are aliquoted and stored at -20°C. Even at -20°C some have a relatively brief shelf life and need to be replaced after a few months. A technique which would enable such enzymes and other molecular biology reagents to be transported and stored at room temperature or above with a long shelf life would be of considerable value.

SUBSTITUTE SHEET

We have found that all restriction enzymes, even the most unstable, can be dried without loss of activity. In the dried form these preparations show very surprising stability and can be kept at up to +55°C for prolonged periods with no detectable loss of activity.

Other sugars and sugar derivatives in contrast have no preservative effect on the stability of these enzymes. Drying in the presence of effective preservatives such as lactitol enables entirely new techniques for using restriction enzymes to be developed. In one such process, the enzymes are dried in the plastic wells of multi-well format microtitration plate (e.g. one with 96 wells). For use, the DNA to be cut is added to the dry well containing the appropriate enzyme in dried form. The water in which the DNA is dissolved, acts as a solvent for the dried enzymes which immediately regain full activity and cut the DNA in a manner identical with the activity of fresh enzyme. This simple one step process should be compared with existing techniques which require that frozen tubes of enzyme be removed from the freezer, aliquoted on ice, diluted in special buffers, added to fresh tubes to which the DNA is subsequently added and then incubated; a more labour intensive operation. For certain applications such as Restriction Fragment Length Polymorphism (RFLP) analysis, a collection of several enzymes is used to cut the same DNA and the pattern of fragments produced by each enzyme is analysed by g l

SUBSTITUTE SHEET

electrophoresis. This technique can be greatly simplified by drying the requisite panel of enzymes in rows of 96 well plates. Thus up to 8 rows of 12 enzymes (or 12 rows of 8 enzymes) can be stored in a single plate. In use, multiple tip pipettes are used to add DNA simultaneously to a whole row of enzymes and all the digests are performed simultaneously with only one manoeuvre.

Dried plates of this type are conveniently stored on a laboratory shelf and do not require either freezing or refrigeration. They are entirely stable for very prolonged periods. Accelerated ageing tests indicate a shelf life of at least 12 months at room temperature. It is probable that the life span is essentially indefinite.

Remarkably, certain sugars which have been claimed by others to have a preservative function are not useful and in some cases positively harmful to the preservation of the biological function of proteins. Thus lactose, and especially palatinose, accelerate the irreversible photobleaching of R-phycoerythrin in solution and completely fail to preserve the activity of restriction enzymes. Sucrose preserves the biological activity of restriction enzymes immediately after drying, but on storage at 37°C progressive damage occurs to the enzymes and after 5 weeks at this temperature, all activity is

SUBSTITUTE SHEET

lost. In contrast trehalose, the disaccharide sugar alcohols lactitol (galactopyranosyl-sorbitol), glucopyranosyl mannitol and glucopyranosyl sorbitol give good preservation of activity. With these sugar alcohols this stabilizing effect is apparently indefinite at elevated temperatures. This is also true of the non-reducing higher sugars such as the trisaccharides raffinose and melezitose. The reducing trisaccharide maltotriose shows no preservative activity on storage at 37°C for 2 weeks. The non-reducing tetrasaccharide stachyose is a good preservative.

In essence, the ability of added carbohydrate to preserve protein stability at elevated temperatures (ambient and above) correlates completely with the absence of a reducing function in the carbohydrate. This can however be an indirect function of the stability of the disaccharide bond. Thus sucrose is a good stabilizing agent short term but within a few weeks at 37°C the protein is destroyed, presumably because of chemical breakdown of the glycosidic bond in sucrose (one of the highest bond energies in biochemistry) which causes secondary chemical damage to the macromolecules.

Absence of a reducing group is not sufficient to guarantee stabilizing ability. Thus the monosaccharide sugar alcohols galactitol, mannitol and erythritol are not satisfactory protective agents. This is not due to

SUBSTITUTE SHEET

the fact that a ring structure is required for stabilizing activity. Sorbitol (glucitol), a straight chain non-reducing monosaccharide alcohol, does have some limited activity as a stabilizing agent while myo-inositol, a non-reducing 6 carbon ring compound is without stabilizing activity.

Of particular importance is the observation that the compounds which stabilize proteins during drying also protect proteins against free radical damage in solution. While we do not wish to be bound by theory, we speculate that this effect is not due to the molecules acting as free radical scavengers because the effective compounds are chemically the most non-reactive and therefore most unlikely to compete for free-radical interaction. It seems more likely that these carbohydrate molecules are able to bind to proteins in solution and act as a buffer between the protein and free radicals produced in the surrounding solvent. Being much larger than water molecules and therefore possessing a much higher moment of inertia, it seems likely that the short lived free radicals cannot displace hydrogen bonded sugar molecules (and damage the protein) within their short life-times. The disaccharide palatinose (isomaltulose) is actively destructive in this test thus showing that single examples of a stabilizing effect of a particular sugar are valuable in predicting its usefulness as a protective agent for biological molecules.

SUBSTITUTE SHEET

W also believe that the ability of effector molecules to coat proteins and other biological molecules with an amorphous phase of hydrogen-bonded carbohydrate is an important element in their protective effect. This is exemplified by experiments in which the fluorescent protein phycoerythrin was dried from 0.6M trehalose solution under conditions in which part of the solidifying mixture crystallised and part solidified as an amorphous glass. Only in the latter phase was the fluorescence of RP-E preserved. Similar data was obtained in drying agarose gels. Where trehalose was used as a preservative, excellent preservation of the agarose structure, as shown by the ability of the dried gel to rehydrate to its original volume, was obtained where amorphous glass transformation occurred. If the trehalose crystallised during drying, the agarose structure was not preserved and the gels in the area in which crystals formed did not reconstitute on addition of water. When agarose gels were dried in the presence of a sugar derivative which always formed an amorphous glass, such as lactitol, the gels showed uniform and complete preservation of structure and function.

The preservative properties of the non-reducing oligosaccharides and oligosaccharide alcohols were not confined to air drying or high temperature storage, nor to the preservation of proteins and carbohydrates. Lyophilised materials also showed excellent protective

SUBSTITUTE SHEET

effects of the same carbohydrates as detailed above. In addition, lipids, specially lipids dispersed as emulsions or liposomes in aqueous buffers could be dried from the same carbohydrate solutions with complete preservation of their properties.

According to the present invention, we thus provide a method of stabilizing delicate biological materials such as proteins, lipids, antibodies etc. for long-term storage under dry conditions, by drying a mutual solution of the biological material and a stabilizing carbohydrate selected from non-reducing sugar alcohol glycosides and non-reducing oligosaccharides having at least three saccharide groups.

We also provide stabilized biological substances formed by drying a solution of the substance in the presence of one of said stabilizing carbohydrates. The invention can thus be seen as having the same applicability as our earlier trehalose process.

We also provide stabilized solutions/suspensions in water.

The following Examples illustrate the invention.

Exempl 1

Th typical restriction endonuclease Pst I is commonly used for molecular cloning and gives a v ry

SUBSTITUTE SHEET

characteristic pattern of phag lambda DNA fragment lengths in electrophoresis gels.

The enzyme was diluted in sterile 96-well U-bottom d polystyrene or PVC plastic plates. These were air dried in a laminar flow hood for 12-16 hours at room temperature (about 18-25°C) and stored at room temperature, 37°C or 55°C for different periods of time for accelerated ageing studies. The drying mixture consisted of three units of the enzyme in 0.45 or 0.6M trehalose or other sugar or sugar derivative in the presence of a proprietary buffer at 10µl/well. The single buffer enabled cutting of DNA with all restriction enzymes and contained a molecular "water pump" component (see PCT/GB89/00093).

<u>Buffer:</u>	33mM	Tris acetate buffer (pH7.4)
	66mM	Sodium acetate
	1mM	Magnesium acetate
	1mM	Dithiotreitol
	3mM	Spermidine
	0.1mg/ml	Bovine Plasma Albumin

Results

Th activity of fresh and dried enzyme was determined by assaying their ability to produce the characteristic pattern of bands from 0.3 µg of lambda

SUBSTITUTE SHEET

bact riophage DNA digested at 37°C for 30 mins and separated by electrophoresis on 1% agarose gels. Enzymes showed no loss of activity after 10 weeks at 55°C. To confirm full retention of enzymatic activity one experiment was also done in which limiting amounts of enzymes were titrated out from 5 units to 0.5 units per well and used to cut 0.3 µg lambda DNA. Trehalose-dried titrated enzymes were compared with titrations of fresh enzymes. After storage for 2 weeks at 37°C the enzyme titred identically to fresh enzyme showing no loss of activity on drying and storage.

Since it is well known that long term damage to macromolecules dried in sugars can result from chemical attack on the macromolecule by the reducing groups in certain mono and disaccharides, these experiments were repeated with several alternative sugars and their non reducing alcohol derivatives. Pst I was dried as above and also in buffers containing a large number of other sugars and chemical derivatives of common sugars. Trehalose again showed 100% preservation (Table 1a and b).

SUBSTITUTE SHEET

Table Ia: Storage of dried Pst I in Different Carbohydrates

CARBOHYDRATE	CHEMICAL NAME	RED	TEMP	TIME	PRESERVATION
Glucose	α -D-glucopyranose	+	37°	1	+
Galactos	α -D-galactopyranose	+	"	14	-
Mannos	α -D-mannopyranose	+	"	1	-
Sorbitol	D-glucitol	-	"	1	-
			"	1	++
			"	14	+
Galactitol	6 carbon alcohol	-	"	70	-
Mannitol	"	-	"	1	-
Erythritol	4 carbon alcohol	-	"	1	-
Myo-Inositol	(1 α ,2 α ,3 α ,4 β ,5 α ,6 β)cyclohexanehexol	-	"	14	-
Thritol	4 carbon alcohol	-	"	14	-
Trehalose	α -D-glucopyranosyl- α -D-glucopyranoside	-	"	14	-
			"	14	+++
			"	14	+++
			"	70	+++
Sucrose	8-D-fructofuranosyl- α -D-glucopyranoside	-	55°	70	+++
			37°	14	+++
Maltose	4-O- α -D-glucopyranosyl-D-glucose	+	"	35	-
			"	14	-
Lactos	4-O- β -D-galactopyranosyl-glucopyranose	+	"	14	-
Palatinose	α -D-glucopyranosyl-1,6-fructose	+	"	14	-
Lactulose	4-O- β -D-galactopyranosyl-D-fructose	+	"	14	-
Maltitol	4-O- β -D-glucopyranosyl-D-glucitol	-	"	14	+++
			"	70	+++
Lactitol	4-O- β -D-galactopyranosyl-D-glucitol	-	"	14	+++
			"	70	+++
			55°	70	+++

SUBSTITUTE SHEET

Table 1b: Storage of Dried Pst I in Various Carbohydrates

CARBOHYDRATE	CHEMICAL NAME	RED	TEMP	TIME	PRESERVATION
Palatinit	1:1 mixture glucopyranosyl -1→6-mannitol+1,6-sorbitol	-	37°	14	+++
Palatinit isomer 1	α-D-glucopyranosyl-1→6- mannitol	-	37°	70	+
Palatinit isomer 2	α-D-glucopyranosyl-1→6- sorbitol	-	"	14	+++
Maltotriose	O-α-D-glucopyranosyl(1→4)- O-α-D-glucopyranosyl(1→4)- D-glucose	+	"	70	+
Raffinos	β-D-fructofuranosyl-O-α-D- galactopyranosyl-(1→6)-α- glucopyranoside	-	"	14	+++
Melezit se	O-α-D-glucopyranosyl-(1→3)- β-D-fructofuranosyl-α-D- glucopyranoside	-	"	14	+++
Stachyose	β-D-fructofuranosyl-O-α-D- D-galactopyranosyl-(1→6)- β-D-fructofuranosyl-α-D- glucopyranoside	-	"	14	+++
Methyl glucoside	methyl-α-D-glucopyranoside	-	"	14	-
M thyl mannoside	methyl-α-D-mannopyranoside	-	"	14	-
Lactulos		+	"	14	+
				35	-

RED = reducing sugar

SUBSTITUTE SHEET

Neither palatinose nor lactose preserved any enzymatic activity. When lactitol was substituted for palatinose or lactose, activity was preserved at all temperatures in a manner equivalent to trehalose i.e. all enzymes were fully active after 10 weeks at 55°C.

Sugar alcohols of the reducing disaccharides gave much better results than the parent disaccharide suggesting that complete lack of reducing activity plus lower-energy glycosidic linkages which resist chemical attack are an important feature of trehalose and other potential preservatives. Interestingly, sugar alcohols of the monosaccharides were poor preservatives except for sorbitol which was only modestly effective. With the higher sugars, again only the non-reducing trisaccharides and the non-reducing tetrasaccharide stachyose were effective.

Example 2

The fluorescent phycobiliprotein R-phycoerythrin is a 240 KD protein extracted from red algae. It has a fluorescence efficiency of 0.98 and emits orange-red light when illuminated with 490 nm blue light. The molecule is very susceptible to desiccation damage and cannot even be lyophilised without collapse of molecular

SUBSTITUTE SHEET

structure and loss of fluorescence. R-PE is also denatured in solution by excitation light in the UV-Blue band of the spectrum. This denaturation is mediated by free radicals created in the solvent water by the irradiation. Free radical damage is also reduced in a dose dependent manner by added trehalose and certain other sugars and sugar alcohols but is, unexpectedly, greatly accelerated by other sugars such as palatinose (Table II)

Table II

Carbohydrate	% loss of fluorescence of R-PE solutions	
	5 min	20 min
trehalose	9%	48%
isomaltulose	93%	100%
lactitol	13%	47%
glucopyranosyl- mannitol (GPM)	21%	47%
glucopyranosyl- sorbitol (GPS)	7%	43%
nil	17%	55%

SUBSTITUTE SHEET

From these results it will be seen that while the two reduced forms of isomaltulose are effective in reducing the loss of fluorescence, isomaltulose itself exhibited a disastrous effect on the fluorescence.

The same panel of sugars and derivatives was used to dry and store a variety of other biological molecules at various temperatures, with essentially similar results.

Example 3

An immunoconjugate produced by covalent bonding of the $F(ab)_2$ fragment of rabbit antibodies against rat immunoglobulin to peroxidase obtained from horseradish.

The conjugate was dried in various sugars and alcohols at a concentration of 15 microgram per ml and 10% sugar by weight and stored for 2 or 4 weeks at 37 or 55°C and titrated for its activity on plates coated with a rat monoclonal antibody.

In trehalose no activity was lost at all. The average values for other sugars and alcohols at 4 weeks were:

GPM	= 86%	; maltitol	67%;	maltose	49%
lactitol	= 86%	; palatinit	68%;	GPS	57%
glucos	= 12%	; no sugar	0%;		

SUBSTITUTE SHEET

Example 4

Agarose gels 7.5 cm x 10 cm x 0.6 cm were poured from molten agarose (1% w/v) containing 15% by weight of various sugars. The gels were dried and stored at various temperatures. When rehydrated, the % regain of wet volume was measured and the gels were used in electrophoresis to separate the Hind III fragments of phage Lambda DNA as a check on function.

Results


All gels failed to rehydrate and to separate DNA after storage at 55°C which seems to be above the re-melting temperature of agarose. At 37°C lactitol and trehalose were perfect from > 12 weeks. Also perfect were GPs and GPM. Palatinose failed after 2 weeks, glucose, mannitol and sorbitol failed after 1 week.

SUBSTITUTE SHEET

CLAIMS:-

1. A method of preserving delicate biological substances or organic compounds (a) in a dry state and/or (b) at elevated temperatures and/or (c) under irradiation comprising incorporating in a system containing the said substances or compounds a sugar or a sugar derivative selected from a (i) non-reducing glycoside of a polyhydroxy compound selected from sugar alcohols and other straight chain polyalcohols, or (ii) a non-reducing oligosaccharide selected from raffinose, stachyose and melezitose.
2. A method according to claim 1 wherein the glycoside is a monoglycoside.
3. A method according to claim 2 wherein the glycoside is obtained by reduction of disaccharide.
4. A method according to claim 1 wherein the glycoside comprises a glucoside or galactoside.
5. A method according to claim 4 when the glycoside is a glycoside of sorbitol or mannitol.
6. A method according to claim 1 wherein the glycoside comprises maltitol; lactitol or one or both isomers of palatinit, the sugar alcohol derived from isomaltulose.

SUBSTITUTE SHEET

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.C1.5 C 12 N 9/96 A 61 K 47/26		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.C1.5	C 12 N A 61 K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	Patent Abstracts of Japan, C-Field, 17 April 1987, & JP, A, 61260882 (WAKAMOTO PHARMACEUT CO. LTD) 19 November 1986, see abstract ---	1-6
X	Patent Abstracts of Japan, 13 November 1986, & JP, A, 61139384 (NIPPON SHINYAKU CO. LTD) 26 June 1986 ---	1-6
X	Chemical Abstracts, vol. 95, 1981, (Columbus, Ohio, US), Ooizumi Tooru et al.: "Quantitative aspect for protective effect of sugar and sugar alcohol against denaturation of fish myofibrils", see page 517, abstract no. 185699n, & Nippon Suisan Gakkaishi 1981, 47(7), 901-8 --- -/-	1-6
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family.</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
19-07-1991	09. 09. 91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE		
Mme Dagmar FRANZ		

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	Chemical Abstracts, vol. 110, 1989, (Columbus, Ohio, US), K. Gekko et al.: "Amino acid solubility and protein stability in aqueous maltitol solutions", see page 292, abstract 110409n, & Agric. Biol. Chem. 1989, 53(1), 89-95 ---	1-6
X	FR,M, 7773 (INSTITUT MERIEUX) 23 March 1970, see page 1, line 36 - page 2, line 6; claim 1 ---	1
X	Japanese Patents Gazette, week 8538, 1985, no. J60149-972A (IATRON LABORATORIES), see abstract ---	1
X	Japanese Patents Gazette, week 8825, 1988, no. J63106-562A (GREEN CROSS), see abstract -----	1

CORRECTED VERSION

ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.

GB 9100759

SA 47465

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on 15/10/91
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR-M- 7773	23-03-70	None	

EPO FORM P0479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82